

**TKN/KS/16/5853**

**Bachelor of Science (B.Sc.) Semester—IV (C.B.S.)  
Examination**

**MATHEMATICS**

**Paper—I**

**(M<sub>7</sub>—Partial Differential Equation and Calculus of  
Variation)**

Time : Three Hours]

[Maximum Marks : 60

- N.B. :—** (1) Solve all the **FIVE** questions.  
(2) All questions carry equal marks.  
(3) Questions **1** to **4** have an alternative.  
Solve each question in full or its alternative  
in full.

**UNIT—I**

1. (A) Find the integral curves of the equations :

$$\frac{dx}{y(x+y)-bz} = \frac{dy}{x(x+y)+bz} = \frac{dz}{z(x+y)}. \quad 6$$

- (B) Verify the equation :

$$x(y^2 - z^2) dx + y(x^2 - z^2) dy - z(y^2 - x^2) dz = 0$$

is integrable and solve it. 6

**OR**

- (C) Verify that the equation :

$$(y^2 + yz) dx + (z^2 + xz) dy + (y^2 - xy) dz = 0$$

is integrable and find its solution. 6

(B) Find P.I. of

$$(D^2 - 6DD' + 9D'^2)z = 12x^2 + 36xy$$

$$\text{where } D = \frac{\partial}{\partial x}, D' = \frac{\partial}{\partial y}. \quad 6$$

**OR**

(C) Solve :

$$\frac{\partial^2 z}{\partial x^2} - \frac{\partial^2 z}{\partial x \partial y} + \frac{\partial z}{\partial y} - z = \cos(x + 2y). \quad 6$$

(D) Solve :

$$x^2 \frac{\partial^2 z}{\partial x^2} - y^2 \frac{\partial^2 z}{\partial y^2} = x^2 y, \text{ by using } x = e^u \text{ and } y = e^v. \quad 6$$

#### UNIT—IV

4. (A) Prove a necessary condition for the functional

$$I[y(x)] = \int_{x_0}^{x_1} F(x, y, y') dx \text{ to be an extremum is that}$$

$$\frac{\partial F}{\partial y} - \frac{d}{dx} \left( \frac{\partial F}{\partial y'} \right) = 0. \quad 6$$

(B) Find the shortest curve joining the two points  $(x_1, y_1)$  and  $(x_2, y_2)$  by using the functional

$$I[y(x)] = \int_{x_1}^{x_2} \sqrt{1 + (y')^2} dx$$

with  $y(x_1) = y_1$  and  $y(x_2) = y_2$ . 6

**OR**

(C) Find the extremum for the functional

$$I[y(x)] = \int_0^\pi (16y^2 - y'^2 + x^2) dx;$$

$$y(0) = y(\pi) = 0, y'(0) = y'(\pi) = 1. \quad 6$$

(D) Write Euler's-Ostrogradsky equation for the

$$\text{functional } I[z(x, y)] = \iint_D \left[ \left( \frac{\partial z}{\partial x} \right)^2 + \left( \frac{\partial z}{\partial y} \right)^2 \right] dx dy.$$

6

#### UNIT—V

5. (A) Find the integral curves of

$$\frac{dx}{1} = \frac{dy}{-2} = \frac{dz}{3x^2 \sin(2x + y)}. \quad 1\frac{1}{2}$$

(B) Form a partial differential equation by eliminating arbitrary constants from the equation

$$z = (x + a)(y + b). \quad 1\frac{1}{2}$$

(C) Find the complete integral of  $pq = 1$ , by Charpit's method. 1\frac{1}{2}

(D) Write the Jacobi's auxiliary equation for  $p^2x + q^2y = z$ . 1\frac{1}{2}

- (D) Prove a necessary and sufficient condition that there exists between two functions  $u(x, y)$  and  $v(x, y)$  a relation  $F(u, v) = 0$ , not involving  $x$  or  $y$  explicitly is that  $\frac{\partial(u, v)}{\partial(x, y)} = 0$ . 6

### UNIT—II

2. (A) Find the general solution of the partial differential equation :

$$z(xp - yq) = y^2 - x^2 \quad 6$$

- (B) Find the integral surface of the partial differential equation  $x^2p + y^2q + z^2 = 0$  through the curve  $xy = x + y, z = 1$ . 6

### OR

- (C) Using Charpit's method, find the complete integral of the partial differential equation :

$$p = (z + qy)^2. \quad 6$$

- (D) Show that a complete integral of  $f\left(\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial u}{\partial z}\right) = 0$

is  $u = ax + by + \phi(a, b)z + c$ , where  $a, b, c$  are arbitrary constants and  $f(a, b, \phi) = 0$ . Further find also the complete integral of

$$\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} = \frac{\partial u}{\partial x} \cdot \frac{\partial u}{\partial y} \cdot \frac{\partial u}{\partial z}. \quad 6$$

### UNIT—III

3. (A) Solve :

$$(D^2 + DD' - 6D'^2)z = y \sin x$$

$$\text{where } D = \frac{\partial}{\partial x}, D' = \frac{\partial}{\partial y}. \quad 6$$

- (E) Solve  $\frac{\partial^2 z}{\partial x \partial y} = 2x + 2y$  by integrating with respect to  $x$  and  $y$ . 1½

- (F) Find P.I. of  $(D^2 - 4DD' + 4D'^2)z = e^{2x+y}$ . 1½

- (G) Let  $I[y(x)] = \int_0^1 [y(x)]^2 dx$  be a functional.

$$\text{If } y(x) = \sqrt{1+x^2} \text{ then find } I[y(x)]. \quad 1½$$

- (H) Find the distance of order zero between the functions  $y = x^2$  and  $y = x$  on the interval  $[0, 1]$ . 1½