

Bachelor of Science (B.Sc.) Semester-IV (C.B.S.) Examination**MATHEMATICS****Paper—I****(M, Partial Differential Equation and Calculus of Variation)**

Time : Three Hours]

[Maximum Marks : 60]

Note :— (1) Solve all the **FIVE** questions.

(2) All questions carry equal marks.

(3) Question Nos. **1** to **4** have an alternative. Solve each question in full or its alternative in full.**UNIT—I**1. (A) Find the integral curves of the equations :

$$\frac{dx}{cy - bz} = \frac{dy}{az - cx} = \frac{dz}{bx - ay}$$

and show that they are circles.

6

(B) Verify the Pfaffian differential equation :

$$(y^2 + z^2)dx + xydy + xzdz = 0$$

is integrable and solve it.

6

OR(C) Verify the equation :

$$z(z + y^2)dx + z(z + x^2)dy - xy(x + y)dz = 0$$

is integrable and find its primitive.

6

(D) Prove a necessary and sufficient condition that there exists between two functions $u(x, y)$ and $v(x, y)$ a relation $F(u, v) = 0$ not involving x or y explicitly is that :

$$\frac{\partial(u, v)}{\partial(x, y)} = 0.$$

6

UNIT—II2. (A) Find the general integral of the linear partial differential equation :

$$z(xp - yq) = y^2 - x^2.$$

6

(B) Show that the equations :

$$xp - yq = x \text{ and } x^2p + q = xz$$

are compatible and hence find their solution.

6

OR

(C) Find the complete integral of the partial differential equation :

$$p^2q^2 + x^2y^2 = x^2q^2(x^2 + y^2)$$

by using Charpit's method.

6

(D) Show that a complete integral of the equation $f\left(\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial u}{\partial z}\right) = 0$ is $u = ax + by + \theta(a, b)z + c$ where a, b, c are arbitrary constants and $f(a, b, \theta) = 0$. Hence find complete integral of the equation :

$$\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} = \frac{\partial u}{\partial x} \cdot \frac{\partial u}{\partial y} \cdot \frac{\partial u}{\partial z}.$$

6

UNIT—III

3. (A) Solve :

$$\frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = -4\pi(x^2 + y^2).$$

6

(B) Solve : $(D^2 - 3DD' + 2D'^2)z = \sin(x - 2y)$

where $D = \frac{\partial}{\partial x}$, $D' = \frac{\partial}{\partial y}$.

6

OR

(C) Solve :

$$\frac{\partial^2 z}{\partial x^2} - \frac{\partial^2 z}{\partial x \partial y} - 2 \frac{\partial^2 z}{\partial y^2} + 2 \frac{\partial z}{\partial y} + 2 \frac{\partial z}{\partial x} = \sin(2x + y).$$

6

(D) Solve :

$$x^2 \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y^2 \frac{\partial^2 z}{\partial y^2} = x^m y^n$$

by using $x = e^a$ and $y = e^b$.

6

UNIT—IV

4. (A) Prove the necessary condition for an extremum of a functional

$$I[y(x)] = \int_{x_0}^{x_1} F(x, y, y') dx \quad \text{where function } F \text{ and its partial derivatives with respect to}$$

x, y and y' upto second order are continuous in the domain D of a plane is

$$F_y - \frac{d}{dx} F_{y'} = 0.$$

6

(B) Find the extremal of the functional :

$$I[y(x)] = \int_0^{\pi/2} (y'^2 - y^2) dx \text{ that satisfies the boundary conditions } y(0) = 0, y(\pi/2) = 1. \quad 6$$

OR

(C) Find the extremal of the functional :

$$I[y(x)] = \int_{x_0}^{x_1} [(y'')^2 - 2(y')^2 + y^2 - 2y \sin x] dx. \quad 6$$

(D) Find Euler-Ostrogradsky equation for the functional :

$$I[u(x, y, z)] = \iiint_D \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 + \left(\frac{\partial u}{\partial z} \right)^2 + 2uf \right] dx dy dz. \quad 6$$

5. (A) Form a partial differential equation by eliminating arbitrary constants from the equation :

$$z = (x + a)(y + b). \quad 1\frac{1}{2}$$

(B) Eliminate the arbitrary function f from the equation $z = xy + f(x^2 + y^2)$ and obtain a partial differential equation. $1\frac{1}{2}$

(C) Find the general integral of the linear partial differential equation $p + q = z. \quad 1\frac{1}{2}$

(D) Show that the equations $f(x, y, p, q) = 0$ and $g(x, y, p, q) = 0$ are compatible if $J_{xp} + J_{yq} = 0. \quad 1\frac{1}{2}$

(E) Solve the partial differential equation $\frac{\partial^2 z}{\partial x \partial y} = 0$ by integration. $1\frac{1}{2}$

(F) Find P.I. of $(D^2 - 2DD' + D'^2)z = \tan(x + y). \quad 1\frac{1}{2}$

(G) Find the distance of order zero between the functions

$$y = xe^{-x} \text{ and } y = 0 \text{ on } [0, 2]. \quad 1\frac{1}{2}$$

(H) Let $I[y(x)] = \int_0^1 [y(x)]^2 dx$ be a functional. If $y(x) = e^x$, then find $I[y(x)]. \quad 1\frac{1}{2}$